

Optimization of the baseline and the parent muon energy for a low energy neutrino factory

Amol Dighe,^{1,*} Srubabati Goswami,^{2,†} and Shamayita Ray^{3,‡}

¹*Tata Institute of Fundamental Research, Homi Bhabha Road, Colaba, Mumbai 400005, India*

²*Physical Research Laboratory, Navrangpura, Ahmedabad 380009, India*

³*Laboratory for Elementary-Particle Physics, Cornell University, Ithaca, NY 14853, USA*

(Dated: October 17, 2011)

We discuss the optimal setup for a low energy neutrino factory in order to achieve a 5σ discovery of a nonzero mixing angle θ_{13} , a nonzero CP phase δ_{CP} , and the mass hierarchy. We explore parent muon energies in the range 5–16 GeV, and baselines in the range 500–5000 km. We present the results in terms of the reach in $\sin^2 \theta_{13}$, emphasizing the dependence of the optimal baseline on the true value of δ_{CP} . We show that the sensitivity of a given setup typically increases with parent muon energy, reaching saturation for higher energies. The saturation energy is larger for longer baselines; we present an estimate of this dependence. In the light of the recent indications of a large θ_{13} , we also determine how these preferences would change if indeed a large θ_{13} is confirmed. In such a case, the baselines ~ 2500 km (~ 1500 km) may be expected to lead to hierarchy determination (δ_{CP} discovery) with the minimum exposure.

I. INTRODUCTION

While the indications of neutrino oscillations first came from the solar and atmospheric neutrino data, the observations from terrestrial experiments have helped provide a firm footing to our knowledge of neutrino masses and mixing. The data from all the neutrino oscillation experiments have established that there are two independent mass squared differences $|\Delta m_{31}^2| \approx 2.35 \times 10^{-3} \text{ eV}^2$ and $\Delta m_{21}^2 \approx 7.58 \times 10^{-5} \text{ eV}^2$, as well as two large mixing angles $\sin^2 \theta_{23} \approx 0.42$ and $\sin^2 \theta_{12} \approx 0.306$ [1, 2]. The third mixing angle θ_{13} is small: at the 3σ level we have an upper bound $\sin^2 \theta_{13} < 0.044(0.035)$ [1]([2]). Recent indications of a nonzero θ_{13} have been obtained at the T2K [3] and MINOS [4] experiments, and now the global fits that incorporate these new data give a non-zero θ_{13} at $\sim 3\sigma$. However, the precise value of the θ_{13} best fit point as well as the significance for $\theta_{13} > 0$ still depend on assumptions on the analysis of data from reactor experiments [2].

The immediate goals for neutrino oscillation experiments are the measurements of (i) the mixing angle θ_{13} , (ii) the CP violating phase δ_{CP} , and (iii) the sign of Δm_{31}^2 , also known as the mass hierarchy. These three quantities, along with the precision measurements of the already known ones, are necessary in order to complete our knowledge of the neutrino mass spectrum.

Among the three quantities mentioned above, θ_{13} is the most important since the determination of the other two depends crucially on the value of this parameter. If $\theta_{13} = 0$, the CP phase is an unphysical quantity, while the determination of mass hierarchy, though possible in principle [5], becomes extremely challenging. If

$\sin^2 2\theta_{13} \gtrsim 0.01$, its measurement will be within the reach of accelerator experiments like T2K, MINOS or NO ν A that use conventional hadron beams, for certain δ_{CP} values. Indeed the recent results [3, 4] indicate that such a measurement may soon be possible. If on the other hand the $\nu_\mu \rightarrow \nu_e$ signals observed here are background or statistical fluctuations, then θ_{13} may be even smaller. Reactor experiments, Double CHOOZ, Reno and Daya-Bay may probe this angle as long as $\sin^2 2\theta_{13} \gtrsim 0.033, 0.018$, and 0.007 respectively [6], independent of the value of δ_{CP} .

The most efficient way of determining mass hierarchy is the observation of the difference in Earth matter effects for the two hierarchies, which is possible if θ_{13} is sufficiently large. Future atmospheric neutrino experiments can achieve this task for $\sin^2 2\theta_{13} \sim 0.04$ at 95% C.L. [7]. For T2K and NO ν A, the sensitivity to hierarchy is possible for $\sin^2 2\theta_{13} > 0.02$ at 90% C.L., albeit only for limited range of values of δ_{CP} [8].

The measurement of δ_{CP} is perhaps the most difficult of the three. Not only does it need a substantial value of θ_{13} , the value of δ_{CP} itself needs to be sufficiently different from zero for a positive signal of CP violation. Planned superbeam experiments, which would use highly intense conventional beams, have a limited sensitivity to δ_{CP} at 90% C.L. and almost no sensitivity at 3σ [8].

While the technology for the conventional beams is well-established, the beam contamination inherent in such beams does not allow measurements accurate to more than a percent level. On the other hand, neutrino production from the decays of muons that are accelerated and stored in a ring (“neutrino factory”), combined with detectors that can identify the charge of leptons produced from the neutrino interactions, has the potential of measuring the quantities of interest even for much smaller θ_{13} values. Even if the conventional beams succeed in an measurement, it is important to confirm such a measurement with another type of source, just like the

*Electronic address: amol@theory.tifr.res.in

†Electronic address: sruba@prl.res.in

‡Electronic address: sr643@cornell.edu

measurements from solar and atmospheric experiments were later confirmed and established by terrestrial neutrino experiments. A neutrino factory is thus a discovery machine in the worst-case scenario (measurements beyond the reach of conventional beams) and a precision machine in favorable scenarios.

In a neutrino factory, accelerated muons are allowed to decay in the long straight sections of a storage ring, resulting in a strong collimated beam. A μ^+ beam decays to give $\bar{\nu}_\mu$ and ν_e . Oscillations of the ν_e to ν_μ produce a μ^- in the detector giving the so called “wrong sign” muon signal, whereas the unoscillated $\bar{\nu}_\mu$ produce the “right sign” or “same sign” muon signal of μ^+ . A detector with a charge identification capability can identify the two different type of signals separately, allowing the determination of $P_{e\mu} \equiv P(\nu_e \rightarrow \nu_\mu)$ and $\bar{P}_{\mu\mu} \equiv P(\bar{\nu}_\mu \rightarrow \bar{\nu}_\mu)$. A μ^- beam would similarly lead to the measurements of $\bar{P}_{e\rightarrow\mu} \equiv P(\bar{\nu}_e \rightarrow \bar{\nu}_\mu)$ and $P_{\mu\mu} \equiv P(\nu_\mu \rightarrow \nu_\mu)$. While the sensitivity stems mainly from the wrong sign muon signal due to the appearance channels $\nu_e \rightarrow \nu_\mu$ and $\bar{\nu}_e \rightarrow \bar{\nu}_\mu$, the disappearance channels $\nu_\mu \rightarrow \nu_\mu$ and $\bar{\nu}_\mu \rightarrow \bar{\nu}_\mu$ also contribute because of the large statistics available in these channels. The sensitivity of a neutrino factory may be up to $\sin^2 2\theta_{13}$ as low as $\sim 10^{-4}$ [9].

Since the appearance probability $P_{e\mu}$ depends on all the three hitherto unknown quantities, the channel $\nu_e \rightarrow \nu_\mu$ in principle contains information on all of them, and hence it has been acclaimed as the golden channel. However this advantage is masked by the fact that the value of δ_{CP} is completely unknown. This gives rise to degenerate solutions making the unambiguous determination of the oscillation parameters an uphill task. An elegant solution was provided by observing that at the distance ~ 7500 km, the golden channel probability $P_{e\mu}$ becomes independent of the CP violating phase irrespective of energy, hierarchy and oscillation parameters [10]. This baseline is the so called “magic” baseline. A neutrino factory is considered most suitable for a magic baseline experiment because of the large flux and the small beam background.

Of course since the oscillation probability at the magic baseline is independent of the CP phase, there is no sensitivity to δ_{CP} . Therefore though the magic baseline experiment is suitable for determination of hierarchy and θ_{13} , one has to consider other shorter baselines for the δ_{CP} determination. Detailed energy-baseline optimization studies by the IDS-NF group propose two magnetized iron neutrino detectors (MIND), one at a distance of 4000 km and another at the distance of 7500 km, with a muon energy of 25 GeV [11]. However this requires high acceleration of the muons, and one has to contend with the $1/r^2$ fall-off of the flux.

In the recent past many authors have investigated the prospect of having a neutrino factory of much lower energy (4 -10 GeV), and hence a shorter baseline. This was termed as the low energy neutrino factory (LENF) [12]. The preferred detectors at these low energies are the magnetized totally active scintillator detectors (TASD) or Liquid Argon detectors that can detect muon change

efficiently. Recently the MIND-type detectors for LENS were also considered [13]. Non-magnetic detectors for LENS has been considered in [14].

The most discussed baseline in the context of LENS is the DUSEL baseline of 1300 km. However recently it was pointed out in [15] that the baseline ~ 2540 km has a special property that in the inverted hierarchy (IH), the golden channel probability $P_{e\mu}$ is independent of the CP phase around 3.3 GeV and hence it can be used for an efficient determination of hierarchy using the $\nu_\mu \rightarrow \nu_e$ channel in superbeams. In a subsequent Letter [16], we showed that for the same baseline, the probability $P_{e\mu}$ in the normal hierarchy (NH) also becomes independent of the CP phase at 1.9 GeV. Therefore we termed this as the “bimagic” baseline and the energies as the magic energies. We also observed that away from the magic energies, the probabilities still depend on δ_{CP} . Therefore an experiment at this baseline would be sensitive to all the three parameters if one uses a broadband neutrino beam from 1–4 GeV as can be obtained from a 5 GeV neutrino factory. A noteworthy point is that the distance 2540 km which was motivated purely from physics considerations in [15] and [16] happens to be close to the Brookhaven–Homestake [17] and CERN–Pyh  salmi [18] baselines ¹.

We note that the LENS was initially motivated to study precision neutrino properties at large θ_{13} . Therefore, with the current indication that θ_{13} is indeed large, an exploration of the potential of a LENS is worthwhile and timely. Since the neutrino-factory technology is still not well-established, the time scale at which these experiments will start is comparatively larger, and therefore its aim should be correspondingly higher – like the measurements of the above quantities of interest to 5σ . This is particularly true for a quantity like the mass hierarchy, which is a binary measurement. A large θ_{13} is also conducive for a measurement of δ_{CP} and it is likely that this aspect may play a decisive role in ascertaining which is the optimal baseline and energy. Of course if θ_{13} happens to be smaller then it needs to be explored what is the optimum baseline and energy for determination of the all the three unknowns – mass hierarchy, θ_{13} and δ_{CP} .

In general, optimization is a complex numerical problem. The parameters involved are energy, baseline as well as the true values of the oscillation parameters. There is also the issue of the optimization with respect to the detector. The dependence of probabilities on the true values of θ_{13} , δ_{CP} and Δm_{31}^2 is beyond the experimental control, so the reach of an experiment has to be assessed in the worst case scenario as far as the values of the mixing parameters are concerned. As we shall show further in this paper the lack of knowledge of true δ_{CP} makes it extremely difficult to zero in on a particular baseline

¹ Potential of baselines close to 2540 km has been studied in [19] in the context of superbeams.

as “the optimal baseline” using the so called green-field approach, where one determines the optimal baseline by a numerical scan of the relevant parameter space.

Optimization studies in the context of low energy neutrino factories have been carried out recently in [13] in the context of a MIND. In this paper we consider a detector that is TAsD. Apart from this, one of the major differences in their analysis and ours is that they present the sensitivity plots in terms of “fraction of δ_{CP} ”, whereas our results are presented for all δ_{CP} values in the range $[0 - 2\pi]$ and our plots reveal the specific range of values of δ_{CP} for which a given baseline is sensitive to a particular quantity.

The plan of the paper goes as follows. In Sec. II we discuss the physics of oscillation probabilities giving us the bimagic condition and the deviations from this condition in the nearby baselines. In Sec. III we discuss the experimental set up and the details of the numerical simulation. We present the results for the baseline optimization in Sec. IV and those for the muon energy optimization in Sec. V. The dependence on the true values of δ_{CP} and $|\Delta m_{31}^2|$ is shown by bands obtained by varying these parameters over their currently allowed ranges. The estimation of an optimal muon energy, given a baseline, is outlined in Sec. VI with a simple approximation. In Sec. VII we discuss the implications of a large measured θ_{13} value for the optimization. Sec. VIII, we summarize our results and comment on future prospects.

II. MAGIC AND BIMAGIC BASELINES

Certain properties of the neutrino flavor conversion probability $P_{e\mu}$ can be useful for an analytical understanding of why certain baselines or parent muon energies should work better than the others. While these have been pointed out earlier [15, 16, 20], we expound on them here in detail, bringing out some of their most important features.

In general the $P_{e\mu}$ oscillation probability can be written as [20]

$$P_{e\mu} = |\cos \theta_{23} A_S e^{i\delta_{\text{CP}}} + \sin \theta_{23} A_A|^2 \quad (1)$$

where A_S is the “solar” amplitude that depends on the solar parameters Δm_{21}^2 and θ_{12} , and A_A is the “atmospheric” amplitude which depends on Δm_{31}^2 and θ_{13} . From the above expression it is evident that the CP violation in neutrino oscillation arises from the interference effects of these two amplitudes. In matter of constant density, the oscillation probability $P_{\nu_e \rightarrow \nu_\mu}$ can be expanded keeping terms upto second order in the small

parameters $\alpha \equiv \Delta m_{21}^2 / \Delta m_{31}^2$ and s_{13} as [21]

$$P_{e\mu} = 4s_{13}^2 s_{23}^2 \frac{\sin^2 [(1 - \hat{A})\Delta]}{(1 - \hat{A})^2} + \alpha^2 \sin^2 2\theta_{12} c_{23}^2 \frac{\sin^2 \hat{A}\Delta}{\hat{A}^2} + 2\alpha s_{13} \sin 2\theta_{12} \sin 2\theta_{23} \cos(\Delta - \delta_{\text{CP}}) \times \frac{\sin \hat{A}\Delta}{\hat{A}} \frac{\sin [(1 - \hat{A})\Delta]}{(1 - \hat{A})}, \quad (2)$$

where $s_{ij} \equiv \sin \theta_{ij}$, $c_{ij} \equiv \cos \theta_{ij}$. Also,

$$\hat{A} \equiv \frac{2\sqrt{2}G_F n_e E_\nu}{\Delta m_{31}^2}, \quad \Delta \equiv \frac{\Delta m_{31}^2 L}{4E_\nu}, \quad (3)$$

where G_F is the Fermi constant and n_e is the electron number density. For neutrinos, the signs of $\hat{A}$ and Δ are positive for normal hierarchy and negative for inverted hierarchy. $\hat{A}$ picks up an extra negative sign for anti-neutrinos. The last term in Eq. (2) corresponds to the interference term from which the CP dependence of the probability originates. This term also mixes the dependence on hierarchy and δ_{CP} , as well as the dependence on θ_{13} and δ_{CP} , leading to a four-fold degeneracy [22]. There is also a degeneracy between $(\theta_{23}, \delta_{\text{CP}})$ and $(\pi/2 - \theta_{23}, \delta_{\text{CP}})$ [23]. Together, this eightfold degeneracy makes the determination of the oscillation parameters ambiguous. It was noticed in [10] that the CP dependence of the probability can be avoided if one has

$$\sin(\hat{A}\Delta) = 0, \quad (4)$$

which corresponds to the vanishing of the solar amplitude A_S and hence the interference term. As a result $P_{e\mu}$ becomes independent of the CP phase δ_{CP} as well as the solar parameters. This condition is obeyed at the so-called magic baseline ($L \sim 7500$ km) for all E_ν and for both the hierarchies.

However, the dependence on the CP phase also vanishes when the atmospheric amplitude A_A vanishes [20], which corresponds to the condition

$$\sin[(1 - \hat{A})\Delta] = 0. \quad (5)$$

However one notes that unlike the magic baseline condition this condition depends on energy as well as hierarchy. Therefore for a particular hierarchy and a particular baseline one can find a set of magic energies where the CP dependence in the probability vanishes. If we consider inverted hierarchy (IH), condition for no δ_{CP} sensitivity (IH-NoCP) can be written as

$$(1 + |\hat{A}|) \cdot |\Delta| = n\pi \quad (6)$$

with the integer $n > 0$. The magic energies are given as

$$E_{\text{magic}}^{\text{IH}} = \frac{1.27|\Delta m_{31}^2|L}{n\pi - 1.27|K|\rho L}. \quad (7)$$

Here the quantity K has been defined such that $K\rho \equiv 2\sqrt{2}G_F n_e$, where $\rho \equiv \rho(L)$ denotes the average matter




FIG. 1: (Color online) Graphically solving the pair of magic conditions. Left panel: $E_{\text{magic}}^{\text{IH}}$ (solid) and $E_{\text{max}}^{\text{NH}}$ (dashed) as a function of L for different n, m . Right panel: $E_{\text{magic}}^{\text{NH}}$ (solid) and $E_{\text{max}}^{\text{IH}}$ (dashed) as a function of L for different n, m . The intersection points of the curves show the baselines and the energies at which the conditions for maximum hierarchy sensitivity are satisfied.

density for the baseline L . For a given baseline L , at these energies the probability $P_{e\mu}$ for IH is independent of the CP phase as well as θ_{13} and only the $\mathcal{O}(\alpha^2)$ term contributes. The hierarchy dependence of the magic energy can be utilized to maximize the hierarchy sensitivity by demanding that $\sin[(1 - \hat{A})\Delta] = \pm 1$ for normal hierarchy (NH) at the same time. We will refer this condition as NH-max. With this condition, the number of events are enhanced due to the first term in Eq. (2). Also the CP dependence is retained in the NH probability. The condition for maxima in NH can be expressed as

$$(1 - |\hat{A}|) \cdot |\Delta| = (m - 1/2)\pi, \quad (8)$$

when m is any integer. This gives

$$E_{\text{max}}^{\text{NH}} = \frac{1.27|\Delta m_{31}^2|L}{(m - 1/2)\pi + 1.27|K|\rho L}. \quad (9)$$

It may be conjectured that we will have maximum hierarchy sensitivity if

$$E_{\text{magic}}^{\text{IH}} = E_{\text{max}}^{\text{NH}}. \quad (10)$$

From the condition in Eq. (10) we arrive at the baseline

$$\rho L (\text{km g/cc}) \approx (n - m + 1/2) \times 16300. \quad (11)$$

Note that the relevant L in Eq. (11) is independent of any oscillation parameters as in the case of the magic baseline. However unlike the magic baseline this condition will be satisfied only for particular values of energy, given by $E_{\text{magic}}^{\text{IH}} = E_{\text{max}}^{\text{NH}}$.

Alternatively if we demand no sensitivity to CP phase in NH (NH-NoCP), we get

$$(1 - |\hat{A}|) \cdot |\Delta| = n\pi, \quad (12)$$

which gives

$$E_{\text{magic}}^{\text{NH}} = \frac{1.27|\Delta m_{31}^2|L}{n\pi + 1.27|K|\rho L}, \quad (13)$$

where n is any non-zero integer. For a given baseline L , at these energies the probability $P_{e\mu}$ for NH is independent of the δ_{CP} as well as θ_{13} , and $P_{e\mu}$ becomes $\mathcal{O}(\alpha^2)$. The condition for maxima for IH (IH-max) gives

$$(1 + |\hat{A}|) \cdot |\Delta| = (m - 1/2)\pi, \quad (14)$$

where m is a positive integer. This gives

$$E_{\text{max}}^{\text{IH}} = \frac{1.27|\Delta m_{31}^2|L}{(m - 1/2)\pi - 1.27|K|\rho L}. \quad (15)$$

Demanding

$$E_{\text{magic}}^{\text{NH}} = E_{\text{max}}^{\text{IH}} \quad (16)$$

we get the same equation for the baseline L , as in Eq. (11), except for an overall negative sign which can be attributed to the different regions of validity for n, m .

Fig. 1 demonstrates the existence of bimagic baselines along with the corresponding magic energies, for $n, m = 1, 2, 3$. The left panel shows the solutions for IH-NoCP and NH-max, while the right panel shows NH-NoCP and IH-max solutions. It is clear from the figure that the baseline $L \sim 2540$ km is the shortest baseline that satisfies both the pairs of conditions simultaneously. It is not a surprise, since the equation for the baseline Eq. (11) is similar for both the pairs of conditions. This baseline is therefore termed as bimagic [16]. For the pair IH-NoCP and NH-max, the solution is obtained from Eq. (11) with $n = m$ and an average matter density of $\rho \sim 3.2$ g/cc. The corresponding magic energies can be obtained from Eq. (7) or Eq. (9) to be 3.3




FIG. 2: (Color online) Conversion probability $P_{e\mu}$ for $L = 1500$ km, 2500 km, 3500 km and 4000 km. The bands correspond to $\delta_{CP} \in [0, 2\pi]$. Other parameters are as given in Eq. (17). The red (thick) line corresponds to $\theta_{13} = 0$.

GeV for $(n, m) = (1, 1)$, 1.4 GeV for $(n, m) = (2, 2)$, 0.89 GeV for $(n, m) = (3, 3)$ and so on. Clearly the magic energies decrease with increasing n, m . Larger values of n, m are less and less practical since the flux at low energies, as well as the efficiency of detection, is typically lower. The solution for the pair NH-NoCP and IH-max is similarly obtained, the magic energies here are 1.97 GeV ($n = 1, m = 2$) and 1.09 GeV ($n = 2, m = 3$).

As can be seen from the figure, the baseline of 6172 km also satisfies the two pairs of conditions. With $\rho(6172 \text{ km}) = 3.955 \text{ g/cc}$, IH-NoCP and NH-max are satisfied for $n - m = 1$; the magic energies in the relevant range are 4.8 GeV for $(n, m) = (2, 1)$ and 2.66 GeV for $(n, m) = (3, 2)$. For NH-NoCP and IH-max, the interesting energy is 3.42 GeV, obtained with $(n, m) = (1, 3)$. This baseline therefore also deserves the title “bimagic”. So do the longer baselines of 8950 km and 10690 km, which are not shown in the figure. However for the purpose of numerical optimization studies in this paper, we restrict ourselves to baselines in the range 500 - 5000 km, since longer baselines imply a lower flux, following the $1/r^2$ behavior.

Even though solving the bimagic conditions given in Eqs. (10) and (16) using the Preliminary Reference Earth Model (PREM) [24] profile we get the exact values of the bimagic baselines with the magic energies, it is observed from Fig. 1 that the values of magic energy for one hierarchy and the maximum energy for the other hierarchy move away from each other rather slowly on either sides of the intersection point, as L is varied. Moreover, cur-

rently there is $\sim 5\%$ error on $|\Delta m_{31}^2|$ at 1σ as well as uncertainties associated with the density profile of earth. So the excellent hierarchy sensitivity of the bimagic baselines, attributed to its bimagic property, is expected to be there even if we move slightly away from these specific baselines. There already exist a few possible baselines of similar magnitudes: (i) Brookhaven to Homestake distance is exactly 2540 km, and (ii) CERN to Pyhäsalmi (proposed site for the LENA detector) distance is 2288 km.

Fig. 2 shows the probability $P_{e\mu}$ for different baselines, for $\sin^2 \theta_{13} = 0.01$. In this and all other plots, we have solved the exact neutrino propagation equation numerically using the PREM profile. The probability $P_{e\mu}$ shown in Fig. 2 shows the presence of the bimagic properties for $L \sim 2500$ km clearly. Around this distance, the maximum of NH appears at the same energy as the minimum of IH and vice versa, enhancing the hierarchy sensitivity. This particular feature is absent in any of the other baselines shown. When one goes to a higher baseline, the following effects occur:

- (i) The flux decreases as $1/r^2$.
- (ii) The amplitude of oscillations in NH increases while that in IH decreases. This would tend to increase the hierarchy sensitivity.
- (iii) The oscillation maxima of IH move faster to higher energies than that of NH, resulting in the maxima of both hierarchies coming closer in energies, which would tend to decrease the hierarchy sensitivity.
- (iv) At a given muon energy, the relevant spectral regions

below the highest probability peak decrease, which also tend to decrease hierarchy sensitivity at that energy. The net result is a subtle combination of all these effects. At the lower end of the parent muon energies E_μ , the bimagic baseline gives the best sensitivity to hierarchy, while at higher E_μ the optimal baseline increases.

The figure also shows the probability for $\theta_{13} = 0$. This is the same for both the hierarchies since in Eq. (2) at $\theta_{13}=0$ only the $\mathcal{O}(\alpha^2)$ term contributes in the leading order which is independent of hierarchy. The distance of the bands from this line gives an direct estimate of θ_{13} discovery potential for the corresponding baseline and hierarchy. The 2500 km plot in Fig. 2 suggests that θ_{13} discovery potential is expected to be good for NH, while it may not be so good for IH as the $P_{e\mu}$ values are lower. The same is true for the longer baselines. The conclusions get reversed for anti-neutrinos, because of the change in sign in $\hat{A}$ in Eq. (2). However for 1500 km, the difference between the probability values for NH and IH is not so large, and hence θ_{13} discovery potential is expected to be similar for both the hierarchies. Of course on top of the probabilities, the cross section and the $1/r^2$ flux dependence also contribute in determining the optimal baseline.

The sensitivity to the CP phase is related to the widths of the bands that represent the variation of the CP phase. The widths of the $P_{e\mu}$ bands seem to increase with increasing baselines. On the other hand, the fluxes fall as $1/r^2$. So the optimal baseline should emerge from a compromise between these two opposing factors.

In this section we have given analytic arguments to motivate the desirable values for the baselines and neutrino energies to determine the sign of mass hierarchy, and detect nonzero values for θ_{13} and δ_{CP} . To choose the optimal baseline for the low energy neutrino factory, one will have to perform a complete numerical study, which we do in the next sections.

III. DETAILS OF NUMERICAL SIMULATIONS

As the detector, we use a 25 kt TASD with an energy threshold of 1 GeV. We choose a typical Neutrino factory setup with 5×10^{21} useful muon decays per year, which is of the same order as in the setup considered in [25, 26]. We consider the running with both the polarities, each for 2.5 years. So we have a neutrino flux consisting of $\bar{\nu}_\mu$ and ν_e when we consider running with μ^+ , while it becomes ν_μ and $\bar{\nu}_e$ when running with negative polarity.

We assume a muon detection efficiency of 94% for energies above 1 GeV, 10% energy resolution for the whole energy range and a background level of 10^{-3} for the $\nu_e \rightarrow \nu_\mu$ and $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ channels. Detection of ν_e or $\bar{\nu}_e$ is not considered in this study, which seems to have a very small effect when the initial flux is as large as above [26]. A 2.5% normalization error and 0.01% calibration error, both for signal and background, have also been taken into account throughout this study. The detector characteris-

tics have been simulated with GLoBES [27]. We also use the prescriptions for priors and marginalization in-built in GLoBES.

Our main goal is to find out the optimal baseline as well as the optimal parent muon energy E_μ for that baseline. In order to optimize the baseline, we choose three representative energies for the parent muon: 5 GeV, 7.5 GeV and 10 GeV, and vary the baseline in the range 500 – 5000 km. For optimizing the parent muon energy E_μ , we choose three representative baselines: 1500 km, 2500 km and 3500 km, and vary E_μ in the range 2 – 16 GeV.

IV. OPTIMIZATION OF THE BASELINE

In this section, we present the results of our baseline optimization for the measurement of the three “performance indicators” [13]: neutrino mass hierarchy, discovery of θ_{13} and δ_{CP} . We have performed the analysis at the parent muon energies of $E_\mu = 5, 7.5$ and 10 GeV. The results at intermediate energies can be extrapolated from the results at these representative parent muon energies. The energy optimization will be presented in the next section.

The main sources of uncertainty in determining the reach of an experiment are the unknown values of δ_{CP} and θ_{13} . We therefore focus on the influence of these two quantities on our results, and keep the true values of other mixing parameters to be fixed at

$$\begin{aligned} \Delta m_{21}^2 &= 7.65 \times 10^{-5} \text{ eV}^2, \quad \sin^2 \theta_{12} = 0.3, \\ |\Delta m_{31}^2| &= 2.4 \times 10^{-3} \text{ eV}^2, \quad \sin^2 \theta_{23} = 0.5. \end{aligned} \quad (17)$$

We also explore the effects of varying $|\Delta m_{31}^2|$ in its current 3σ allowed range. In this section we present three kinds of plots:

- (i) Type-A: plots in the $\sin^2 \theta_{13} - L$ plane for fixed true values of $|\Delta m_{31}^2|$ and δ_{CP} varying in the range $[0, 2\pi]$,
- (ii) Type-B: plots in the $\delta_{\text{CP}} - L$ plane for fixed true values of $\sin^2 \theta_{13}$ and $|\Delta m_{31}^2|$,
- (iii) Type-C: plots in the $\sin^2 \theta_{13} - L$ plane for fixed true values of δ_{CP} and varying $|\Delta m_{31}^2|$ in its current 3σ range.

In the conventional plots the reach for a particular performance indicator is often given in terms of the fraction of δ_{CP} values for which the determination of a quantity is possible. Our Type-A plots show the reach for all possible δ_{CP} values, from which the reach even for the worst-case δ_{CP} values may be inferred. Moreover, from the Type-B plots that show the reach for all δ_{CP} values, one may also trivially infer the fraction of δ_{CP} for which the quantity may be determined.

For reference, in the baseline optimization plots we also show two vertical lines, corresponding to the baselines of 2540 km, the “bimagic” baseline, and 1300 km, the baseline that is perhaps the most studied in the context of the LENF.

A. Hierarchy determination

In order to optimize the baseline for the determination of hierarchy, we assume NH to be the true hierarchy and show the reach of θ_{13} for which the wrong hierarchy (IH) can be excluded to 5σ , as a function of the baseline. Note that the determination of hierarchy is a binary measurement, and hence a 5σ determination is absolutely necessary before claiming a positive identification of this quantity.

Figure 3 shows the hierarchy sensitivity with the stated experimental setup. In the top panel we present Type-A plots, where true values of $\sin^2 \theta_{13}$ are plotted along the vertical axis. True values of all other parameters, except δ_{CP} , are set to values stated in Eq. (17). To generate the bands in this top panel, $\delta_{\text{CP}}(\text{true})$ is varied over the complete range of $[0, 2\pi]$. For each set of chosen values of mixing parameters and chosen baseline, χ^2_{min} is obtained by marginalizing over all parameters with the wrong hierarchy. We have taken 4% error on each of Δm_{21}^2 and θ_{12} and 5% on θ_{23} and $|\Delta m_{31}^2|$. A 2% error has also been considered on the earth matter profile and marginalized over.

The figures in the top panel of Fig. 3 show that the baseline optimization depends on the actual δ_{CP} value as well as the parent muon energy E_μ . The widths of the bands, which span almost an order of magnitude in $\sin^2 \theta_{13}$, are mainly due to the variation in δ_{CP} , while the dependence on E_μ may be discerned from the variation across the three plots in the panel. For each baseline, the upper edge of the band gives the lowest value of θ_{13} for which hierarchy can be determined at 5σ in the most conservative case, i.e. irrespective of what the true value of δ_{CP} is.

The leftmost top panel shows that for $E_\mu = 5$ GeV, the reach of the experiment is optimal at $L \sim 2500$ km for most of the δ_{CP} values, however the actual reach depends strongly on the true δ_{CP} . At the optimal baseline the reach in $\sin^2 \theta_{13}$ varies over an order of magnitude, from $\sim 10^{-4}$ to 10^{-3} . The maximum reach around this baseline is for $\delta_{\text{CP}}(\text{true}) = 3\pi/4$, which is consistent with the analytic estimate obtained in [16]. For increasing parent muon energies, the optimal baseline typically increases. For longer baselines the maximum of NH and minimum of IH shifts to higher energies. Although the condition $E_{\text{max}}^{\text{NH}} = E_{\text{magic}}^{\text{IH}}$ is not exactly satisfied, the broader width of the oscillation curve for higher baselines contributes to give an enhanced sensitivity. The widths of the bands in the top panel are rather conservative, since they include all true values of δ_{CP} .

In reality there is a unique true value of δ_{CP} , which controls the reach for $\sin^2 \theta_{13}$ for a given baseline. Since this true value is unknown in the analysis it has to be kept free. In order to understand what are the specific values of δ_{CP} for which hierarchy can be determined at a particular baseline, we also present the Type-B plots in the middle panel of Fig. 3, plots for fixed θ_{13} values in the $\delta_{\text{CP}} - L$ plane.

The middle-panel plot for 5 GeV muon energy shows that if $\sin^2 \theta_{13}$ is 0.056, then it is possible to determine hierarchy at 5σ level for all values of δ_{CP} for $L \geq 700$ km. For $\sin^2 \theta_{13} = 0.01$, one needs to go beyond 1000 km if $\delta_{\text{CP}} > 3\pi/2$. For $\sin^2 \theta_{13} = 0.001$, there is some sensitivity to hierarchy for $L \simeq 1100$ km, but only for δ_{CP} close to $\pi/2$. Hierarchy sensitivity at 1300 km for this value of θ_{13} exists only for $\sim 40\%$ of the possible δ_{CP} values. The baseline that offers sensitivity to the largest range of δ_{CP} values at such low θ_{13} is 1800–2500 km. As we go beyond 3000 km the range of δ_{CP} values for which hierarchy can be determined becomes much smaller for $E_\mu = 5$ GeV. However if the energy is increased, then higher baselines can determine hierarchy for a wider range of δ_{CP} values. One striking feature, seen from these plots, is that there is a small range of δ_{CP} values around $3\pi/2$ for which hierarchy cannot be determined by LENF for $\sin^2 \theta_{13} = 0.001$. This can be remedied by a combination with another experiment that is sensitive to the region around $\delta_{\text{CP}} = 3\pi/2$.

Another uncertainty in the hierarchy sensitivity of a given experimental setup can come from $|\Delta m_{31}^2|$, as the magic energies depend on it. To illustrate this dependence, we show Type-C plots in the bottom panel of Fig. 3 for three fixed $\delta_{\text{CP}}(\text{true})$ values: $\delta_{\text{CP}}(\text{true}) = 0$ (no CPV), $\pi/2$ (maximum CPV), and $3\pi/4$ (intermediate CPV). The true value of $|\Delta m_{31}^2|$ is varied over its 3σ range, giving rise to bands with a finite width. Each plot shows the bands for the three representative E_μ values, stated before.

The widths of the bands in the bottom panel of Fig. 3 show that the dependence of hierarchy sensitivity on $|\Delta m_{31}^2|$ is much smaller compared to that on the true value of δ_{CP} . From the left most figure in the panel, we see that for no CPV, the best hierarchy sensitivity is obtained in a rather broad baseline regime 1500 – 3500 km, though for the worst-case values of $|\Delta m_{31}^2|$, the range around 2500 km is preferred, where the error due to $|\Delta m_{31}^2|$ is also small. If CPV is maximum ($\delta_{\text{CP}} = \pi/2$, central panel) or has the chosen value of $3\pi/4$ (right panel), the maximum hierarchy sensitivity is for a comparatively narrower, but still wide, baseline range $\sim 2300 - 3500$ km, near the bimagic baseline. At the worst-case values of $|\Delta m_{31}^2|$, a baseline ~ 2500 km is preferred for $E_\mu = 5$ GeV, while it shifts to ~ 3000 km for higher E_μ values. Among the three δ_{CP} values considered, the sensitivity to hierarchy is observed to be the worst for the scenario with no CPV, however as can be seen from the top panel, different values of δ_{CP} are the worse-case scenarios for different baselines. As the parent muon energy increases, the optimal baseline shifts to higher values as expected. The sensitivity is seen to increase when E_μ increases from 5 to 7.5 GeV, but thereafter it seems to saturate. This feature will be discussed in detail in the next sections.

Our results in this section clearly indicate that if hierarchy sensitivity is considered as the performance indicator, and if $\sin^2 \theta_{13} \lesssim 0.001$ and $E_\mu = 5$ GeV, then




FIG. 3: (Color online) 5σ reach in hierarchy determination for fixed muon energies, as a function of baseline, assuming the true hierarchy to be NH. The top panel gives the reach in $\sin^2 \theta_{13}$, the bands correspond to $\delta_{CP} \in [0, 2\pi]$. Specific values of δ_{CP} are also shown within the band. The middle panel shows the 5σ reach in δ_{CP} for fixed values of $\sin^2 \theta_{13}$. The bottom panel denotes the reach in $\sin^2 \theta_{13}$ for fixed values of δ_{CP} . The bands correspond to the current 3σ range of $|\Delta m_{31}^2|$. All the undisplayed parameters are fixed at values given in Eq. (17). The plots are generated for 2.5 years of running with each muon polarity. For reference, we also show two vertical lines, corresponding to the baselines of 2540 km (the bimagic baseline), and 1300 km.

the hierarchy can be determined for a larger ($\sim 86\%$) fraction of possible δ_{CP} values at 2540 km, the bimagic baseline. This fraction increases for higher E_μ values. For $\sin^2 \theta_{13} > 0.01$, hierarchy determination is possible for all values of δ_{CP} , as long as $E_\mu > 5$ GeV and $L > 1300$ km.

B. θ_{13} discovery

In this section, we apply the same analysis techniques as the last section to the discovery potential of θ_{13} . Note that though a $\sim 2.5\sigma$ evidence for nonzero θ_{13} has recently been claimed by experiments [3, 4] and the global fit to the neutrino mixing parameters [1, 2], the jury is still out on this and it is quite possible that the value of

θ_{13} is much smaller. Also, since we are looking towards a long-term experiment, we should not be satisfied with a 3σ evidence, but should try to gauge the potential of an experiment for a definitive, 5σ discovery. Fig. 4 shows the 5σ reach for θ_{13} discovery for three different parent muon energies, in terms of the three types of plots, Type-A, B and C, as mentioned above.

The widths of the bands in the top panel show that θ_{13} discovery potential of a baseline depends strongly on the true value of δ_{CP} . Indeed for any baseline, the best-case and the worst-case values of δ_{CP} make a difference of almost an order of magnitude in the corresponding reach for $\sin^2 \theta_{13}$. Shorter baselines are seen to be generally preferred for θ_{13} discovery, as compared to those preferred for hierarchy determination. For example, for $E_\mu = 5$ GeV, the optimal baseline is $\sim 800 - 1600$




FIG. 4: (Color online) 5σ reach in θ_{13} discovery for fixed muon energies, as a function of baseline, assuming the true hierarchy to be NH. The same conventions as in Fig. 3 are used. In the middle panel, for $\sin^2 \theta_{13} = 0.001$, the discovery of θ_{13} is possible for regions outside the dashed (violet) contour. For $\sin^2 \theta_{13} = 0.0001$, the discovery of θ_{13} is possible for the area inside the solid (blue) contours.

km, while for higher parent muon energies, it moves to $\sim 1500 - 2500$ km. At all these energies, it is observed that the sensitivity is maximum when $\delta_{CP} \approx \pi/2$. With the worst-case values of δ_{CP} , the optimal baseline stays near 1500 km in the whole energy range.

In the middle panel of Fig. 4 we present the Type-B plots. For $\sin^2 \theta_{13} = 0.001$, the discovery of θ_{13} is possible for regions outside the violet (dashed) contour. Thus for $L < 2100$ km, θ_{13} can be discovered irrespective of δ_{CP} for this value of $\sin^2 \theta_{13}$. Higher θ_{13} values can be discovered in the whole $\delta_{CP} - L$ plane. For $\sin^2 \theta_{13} = 0.0001$, the discovery of θ_{13} is possible for the area inside the solid blue contours i.e. only for shorter baselines and a limited range of δ_{CP} . For higher baselines like $L \sim 3000$ km, θ_{13} discovery is possible only if E_μ is high and δ_{CP} has values close to $\pi/2$ or $7\pi/4$. In general for such small values of $\sin^2 \theta_{13}$, the baseline of ~ 1500 km seems to have θ_{13} sensitivity for a wider range of δ_{CP} values.

In the bottom panel of Fig. 4, we present the Type-C plots for θ_{13} sensitivity. It illustrates the dependence of the optimal baseline on the true value of δ_{CP} : while for $\delta_{CP} = 0$ the optimal baseline seems to be $\sim 1300 - 1500$ km, for $\delta_{CP} = \pi/2$ it is $\sim 1500 - 2000$ km and for $\delta_{CP} = 3\pi/4$, the most efficient baseline is ~ 2500 km.

To summarize, if θ_{13} discovery is chosen as the performance indicator, baselines in the region $1500 - 2000$ km seems to work best, the actual value of the optimal baseline being dependent on the true value of δ_{CP} . However if $\sin^2 \theta_{13} > 0.001$, then a LENF with E_μ in the range 5 - 10 GeV can discover θ_{13} at 5σ for any baseline in the range 500-2500 km for almost all values of true δ_{CP} .




FIG. 5: (Color online) 5σ reach in δ_{CP} discovery for fixed muon energies, as a function of baseline, assuming the true hierarchy to be NH. The same conventions as in Fig. 3 are used. In the middle panel, δ_{CP} discovery is possible for enclosed regions that do not include $\delta_{\text{CP}} = 0$ or π .

C. δ_{CP} discovery

Next we move to the δ_{CP} discovery potential in this section. The top panel of Fig. 5 shows the 5σ reach of $\sin^2 \theta_{13}$ for δ_{CP} discovery, given a true δ_{CP} value. True values of all other parameters have been fixed to values mentioned in Eq. (17), while the true value of δ_{CP} is chosen as $\pi/4$, $\pi/2$, $3\pi/4$, $5\pi/4$, $3\pi/2$ and $7\pi/4$. We do not choose $\delta_{\text{CP}}(\text{true})$ too close to zero or π since the discovery potential is expected to be low. Finally χ^2_{min} is obtained by marginalizing over all parameters except δ_{CP} , for each $\sin^2 \theta_{13}(\text{true})$. Errors on different parameters are taken to be the same as stated in Section IV A.

It is observed that the best-case sensitivity for δ_{CP} is when δ_{CP} is near $\pi/2$ or $3\pi/2$. This is expected since it is at these values that the CP violation is maximum. Typically for all δ_{CP} values and all parent muon energies, the baselines $\sim 800 - 2000$ km seem to be the most efficient. For $\delta_{\text{CP}} = 3\pi/4, 5\pi/4$, on the other hand, the sensitivity

is much lower. Nevertheless, baselines of $\lesssim 2000$ km are preferred.

The middle panel of Fig. 5 gives the sensitivity to δ_{CP} as a function of baseline for three fixed values of $\sin^2 \theta_{13}$ which are 0.056, 0.01 and 0.001 and three values of energy $E_\mu = 5, 7.5$ and 10 GeV. These are representative values and the results for intermediate values of $\sin^2 \theta_{13}$ can be adjudged. The figure shows that for $\sin^2 \theta_{13} > 0.01$, if δ_{CP} lies in the ranges $(0.3 - 0.7)\pi$ or $(1.3 - 1.7)\pi$, it is possible to discover nonzero δ_{CP} at 5σ for all baselines and all energies considered here, though smaller baselines are a bit more efficient. For lower θ_{13} values, only shorter baselines, $L \lesssim 2000$ km, have the possibility of detection of CP violation for most of the δ_{CP} range. Longer baselines need higher muon energies and specific δ_{CP} values near $\pi/2, 3\pi/2$ in order to achieve the task.

The bottom panel of Fig. 5 gives the Type-C plots for δ_{CP} discovery for three fixed values – $\pi/2$, $3\pi/4$, $3\pi/2$. This figure corroborates that the sensitivity to CP vi-

olation discovery is highest for $\delta_{\text{CP}} = 3\pi/2$. The distance at which the best sensitivity is reached is ~ 1700 km (for $\delta_{\text{CP}} = \pi/2$) and ~ 2100 km (for $\delta_{\text{CP}} = 3\pi/2$). For $\delta_{\text{CP}} = 3\pi/4$ the range 1500 - 2500 km have best sensitivity. In general the sensitivity is better at higher energies.

To summarize, the δ_{CP} discovery potential at various baselines depends on the true $\sin^2 \theta_{13}$. For $\sin^2 \theta_{13}$ upto 0.001, δ_{CP} that is not too close to 0 or 2π can be discovered at 5σ by the LENF setup for baselines 500 – 2000 km. In general lower baselines can access a wider range of δ_{CP} values.

V. OPTIMIZING PARENT MUON ENERGY E_μ

As seen in the last section, the optimal values of the baselines depend on the parent muon energies in addition to the true values of neutrino mixing parameters. In this section, we choose three representative values for the baseline – 1500, 2500 and 3500 km – and perform optimization over the parent muon energies. We use three different performance indicators as before: hierarchy determination, θ_{13} discovery and δ_{CP} discovery. The parent muon energy range considered here is 2 – 16 GeV, covering the range of the proposed LENFs. We present only plots of Type-A, since most of the features of the energy dependence may be obtained through interpolation using the representative values $E_\mu = 5, 7.5$ and 10 GeV chosen in the last section.

A. Hierarchy determination

The top panel of Fig. 6 shows the 5σ hierarchy determination range for the three chosen benchmark baselines. For each chosen true value of $\sin^2 \theta_{13}$ and E_μ , we obtain χ_{min}^2 by marginalizing over all parameters assuming IH, while the true hierarchy is taken to be NH. All the parameters, except δ_{CP} , are fixed to the values quoted in Eq. (17) and the band is generated by varying δ_{CP} in the full range $[0, 2\pi]$. Errors on different parameters have been taken to be the same as described in Section IV A.

Similar to the case of baseline optimization in Fig. 3, the large width of the band indicates the strong dependence of hierarchy determination potential for a given experimental setup on the true value of δ_{CP} . As far as the dependence on E_μ is concerned, the sensitivity increases with E_μ at low E_μ values, and saturates at a certain value of E_μ beyond which there is virtually no change in the sensitivity. Let us refer to this energy as the saturation energy, E_μ^{Sat} . It is observed that E_μ^{Sat} increases with increasing baseline, the values being $E_\mu^{\text{Sat}} \sim 5$ GeV for 1500 km, $E_\mu^{\text{Sat}} \sim 7$ GeV for 2500 km and $E_\mu^{\text{Sat}} \sim 10$ GeV for 3500 km. The sensitivity reached at E_μ^{Sat} also increases with the baseline.

The saturation behavior indicates that one will tend to get significantly better sensitivity with increasing E_μ and

baseline till some limit, beyond which the gain may not be worth the increased acceleration required. We observe that E_μ^{Sat} depends significantly on δ_{CP} as well, however this dependence is not a straightforward one. The sensitivity at a baseline of 2500 or 3500 km is always better than that at 1500 km, sometimes by up to an order of magnitude, for energies beyond ~ 6 GeV. The performances at 2500 and 3500 km are comparable: the latter is marginally better, but requires higher E_μ to achieve its full potential.

Thus if hierarchy determination is considered as the performance indicator, then a LENF can determine hierarchy for $\sin^2 \theta_{13}$ as low as 0.002 at a baseline $L \sim 2500$ km for $E_\mu > 5 \text{ GeV}$, for any $\delta_{\text{CP}}(\text{true})$. If true $\sin^2 \theta_{13}$ happens to be large, then shorter baselines and smaller energies would be sufficient.

B. θ_{13} discovery

In this section we present the result of our study of θ_{13} discovery potential as a function of the parent muon energy E_μ , for the three benchmark baselines. The middle panel of Fig. 6 shows the 5σ contours when $\delta_{\text{CP}}(\text{true})$ is varied in $[0, 2\pi]$ and also for seven chosen true values of δ_{CP} . True values of all other parameters other than the displayed ones have been fixed to the values mentioned in Eq. (17), and χ_{min}^2 is obtained by marginalizing over all except θ_{13} , with the errors stated in Section IV A.

These figures also confirm that the discovery potential of any experimental setup depends crucially on the knowledge of δ_{CP} . The sensitivity is the best when $\delta_{\text{CP}} \approx \pi/2$, however the conservative reach for θ_{13} may be up to an order of magnitude worse. Larger baselines require larger muon energies to get an equivalent sensitivity, however with such a larger E_μ , their reach in θ_{13} may be better.

The saturation behavior as in the case of hierarchy determination – i.e. the sensitivity increases with E_μ till a saturation value E_μ^{Sat} – is observed even for θ_{13} discovery for most δ_{CP} values, prominently for baselines of 1500 and 2500 km, but to a smaller extent for 3500 km.

C. δ_{CP} discovery

The analysis of the optimal parent muon energy for δ_{CP} discovery is carried out on the same lines as that in Sec. IV C. The plots in the bottom panel of Fig. 6 show the 5σ discovery potential contours as a function of E_μ , for six chosen true values of δ_{CP} , for each of the three representative baselines.

It is observed that the best sensitivity would be reached when $\delta_{\text{CP}} = \pi/2$. While in this optimistic scenario the sensitivity of a 3500 km baseline with $E_\mu > 8$ GeV will be better, for the worst-case values of δ_{CP} a baseline of 1500




FIG. 6: (Color online) 5σ reach in $\sin^2 \theta_{13}$ for (top panel) hierarchy determination, (middle panel) θ_{13} discovery, and (bottom panel) nonzero δ_{CP} discovery, as a function of E_μ , for fixed δ_{CP} values. NH is taken as the true hierarchy. All parameters except δ_{CP} are fixed at values quoted in Eq. (17). The exposure is taken to be 2.5 years of running with each polarity.

km is preferred, with $E_\mu \gtrsim 5$ GeV where the saturation is reached.

VI. OPTIMAL PARENT MUON ENERGY FOR A GIVEN BASELINE

The two design parameters of an LENF that can in principle be controlled are the baseline and the parent muon energy that determines the neutrino fluxes at the source. Ideally one can look for the optimal combination of both of these, however it may not always be practical. In particular, considerations behind choosing a baseline involve factors like the location suitable for an accelerator and a location where a underground laboratory for neutrino detection can be built. Apart from scientific considerations, this involves geography, economics as well as sociology. A survey of such pairs of locations was recently carried out [13]. Given a baseline, the choice

of the parent muon energy is relatively straightforward, limited mainly by technological considerations. The energy of the muon beam can then be chosen based on the optimality analysis.

As has been noticed multiple times in the previous two sections, the performance at a baseline typically increases with increasing parent muon energy E_μ (in the LENF range), till it saturates at a particular value E_μ^{Sat} . Given that increasing the muon energy is associated with additional costs, a desirable thing to do would be to choose the value of E_μ at or near the value of E_μ^{Sat} . In this section, we obtain the E_μ^{Sat} values for arbitrary baselines using a simple approximation. It turns out that this approximation matches the numerical E_μ^{Sat} values observed in the previous section.

We expect that the main factor influencing the efficiency at a given energy will be the number of wrong-sign muon events. While the actual numbers will depend on the detector characteristics, we estimate this efficiency




FIG. 7: (Color online) The optimal value E_μ^{opt} as a function of the baseline, for normal hierarchy and with $\sin^2 \theta_{13} = 0.001$ and 0.01.

simply through the “quality factor”

$$Q \equiv \frac{\int \Phi_{\nu_e} P_{e\mu} \sigma_{\nu_\mu} dE_{\nu_e}}{\int \Phi_{\nu_e} \sigma_{\nu_\mu} dE_{\nu_e}} \quad (18)$$

where Φ_{ν_e} is the flux of ν_e at the source and σ_{ν_μ} is the total charged-current cross section of ν_μ at the detector. We have normalized the events to a complete $\nu_e \rightarrow \nu_\mu$ conversion. Since the parent muon energy at the source determines the neutrino spectra at the source completely, and the baseline determines the oscillation probability completely (modulo our knowledge about the mixing parameters), the quality factor is known for a energy-baseline combination. The optimal energy E_μ^{opt} is the one for which this quality factor is maximum. There will be a spread in E_μ^{opt} due to the uncertainties in mixing parameters, mainly θ_{13} , $|\Delta m_{31}^2|$ and δ_{CP} .

Fig. 7 shows the E_μ^{opt} values as function of baseline L , for $\sin^2 \theta_{13} = 0.001, 0.01$. The width is obtained due to variation of E_μ^{opt} as $|\Delta m_{31}^2|$ is varied in the current 3σ range and δ_{CP} in $[0, 2\pi]$. This gives a good estimation of what E_μ values will be optimal corresponding to a given baseline, and can work as a rough guideline for determining the optimal range of E_μ , once the baseline L has been determined from other considerations. Then the upper limit can be used to estimate E_μ^{Sat} for a given baseline. Going beyond E_μ^{Sat} would not improve the performance of the setup.

VII. IF θ_{13} IS LARGE

The indications of a large θ_{13} value from the experiments [3, 4] and the global fit to neutrino data [1, 2] is a good news for the measurements of other neutrino parameters too. In particular, it makes it easier to determine the mass hierarchy, and makes the measurement of δ_{CP} possible.

The figures presented in the previous section already give us some idea about what can be the optimal energies and baseline for hierarchy and δ_{CP} for $\sin^2 \theta_{13}$ in the above range.

If $\sin^2 \theta_{13}$ is close to its present best-fit, then the LENF set up considered in this paper can determine hierarchy irrespective of what is the true δ_{CP} for baselines $\gtrsim 1000$ km, with a 2.5 years exposure with each of the polarities and 5×10^{21} useful muon decays, as shown in Figure 3. If true δ_{CP} is not too close to $3\pi/2$ then even baselines $\gtrsim 700$ km would be sufficient for this purpose. The task of δ_{CP} determination will also be easier: δ_{CP} in the range $(0.2 - 0.8)\pi$ and $(1.2 - 1.8)\pi$ can be determined at 5σ for baselines of 500–2000 km for E_μ in the range 5–7 GeV, as can be seen from Figure 5.

We can then be more ambitious and try to optimize the setup in order to get a 5σ determination of hierarchy and δ_{CP} with the minimum exposure. This is shown in Fig. 8, where we take the range of $\sin^2 \theta_{13}$ to be $[0.013, 0.028]$, the 1σ range given in [1], in the prior and then marginalized over θ_{13} .

It is clear from the top panel of the Fig. 8 that for any parent muon energy, the baseline $L \sim 2500$ km is the optimal one for the determination of mass hierarchy. As mentioned before, this is near the bimagic baseline indicated in [16]. For a baseline of 1500 or 3500 km, the exposure needed may be an order of magnitude larger, depending on the δ_{CP} value.

The bottom panel of Fig. 8 shows that for any E_μ a baseline ~ 1300 km will show the best sensitivity for δ_{CP} discovery, for most of the true values of δ_{CP} . However, the “bimagic baseline” 2540 km will have comparable sensitivity with an exposure ~ 1.5 times larger. If $\delta_{\text{CP}}(\text{true})$ happens to be close to $5\pi/4$, ~ 2500 km may show a better sensitivity compared to 1300 km. Note that the exposure needed for δ_{CP} discovery at 5σ is typically one order of magnitude higher than that for hierarchy determination.

VIII. SUMMARY AND CONCLUDING REMARKS

The neutrino mixing angle θ_{13} , the CP-violating phase δ_{CP} and the hierarchy of neutrino mass eigenstates are the three quantities whose measurements are crucial in order to complete and confirm our current picture of neutrino mixing and oscillations. The next generation of long-baseline neutrino experiments therefore justifiably consider these measurements as their primary goals.

In this article, we go beyond the conventional beam experiments and try to optimize the setup for a low-energy neutrino factory (LENF) – where the energy of the parent muon is less than ~ 15 GeV, and consider a magnetized totally-active scintillator detector (TASD) that can identify muon charge, for these measurements. In the worst case scenario i.e θ_{13} turning out to be smaller than the reach of the current reactor and superbeam experiments such setups can be used as discovery machines. On the other hand if θ_{13} is confirmed to be in the currently indicated range then one can use such machines for precision studies with the aim to measure the three quantities at




FIG. 8: (Color online) Exposure required for 5σ hierarchy determination (top panel) and δ_{CP} discovery (bottom panel), if θ_{13} is large. We have taken $\sin^2 \theta_{13} \in [0.013, 0.028]$ [1]. NH is taken as the true hierarchy, and all parameters except δ_{CP} has been fixed at the values stated in Eq. (17).

the 5σ level. In this paper we have focussed on the discovery potential.

We note that some of the parameters that determine the efficiency of a neutrino factory are beyond our control: the probability $P_{e\mu}$ that controls this efficiency depends crucially on θ_{13} and δ_{CP} . However there are two parameters in our control: the baseline and the energy of the parent muon. It is with respect to these parameters that we perform our optimization.

It is of course obvious that the value of θ_{13} would determine how efficient an experiment is. And that is true more or less in a straightforward way. In the absence of matter effects, $P_{e\mu}$ is proportional to θ_{13}^2 . In the presence of matter effects, θ_{13} affects the probability linearly, thus influencing the identification of mass hierarchy. The term in the probability that involves δ_{CP} is also linear in θ_{13} . Thus at shorter baselines (low matter effects), the value of θ_{13} affects the measurement of θ_{13} itself through a quadratic term, while at larger baselines, it affects the probability through a term that contains the product of itself with the matter effects.

In addition to θ_{13} , we find that even the true value of δ_{CP} can have a large impact on the reach of an LNF experiment. Indeed, the reach can change by up to an order of magnitude or more depending on the true value of δ_{CP} . The actual value of $|\Delta m_{31}^2|$ also affects the efficiency, albeit to a smaller extent. Given that the actual value of δ_{CP} is crucial, we present our results in a way that bring out the impact of this parameter. We present three types of plots: (i) Type-A: plots in the $\sin^2 \theta_{13} - L$ plane for fixed values of $|\Delta m_{31}^2|$ and δ_{CP} varying in the

range $[0, 2\pi]$, (ii) Type-B: plots in the $\delta_{CP} - L$ plane for fixed values of $\sin^2 \theta_{13}$ and $|\Delta m_{31}^2|$, (iii) Type-C: plots in the $\sin^2 \theta_{13} - L$ plane for fixed values of δ_{CP} and varying $|\Delta m_{31}^2|$ in its current 3σ allowed range.

Most of the earlier work with the optimization of LNF presented the results in terms of “fraction of δ_{CP} ” where the experiment was successful. However knowing the exact range of δ_{CP} where a certain measurement is possible is crucial, especially when one is thinking about combining results from two complementary experiments: these experiments should be sensitive to complementary ranges of δ_{CP} .

Our detailed observations may be found in the main body of this text. We would like to attract the reader’s attention to some salient features. The following observations refer to 2.5 years of running with each muon polarity, with 5×10^{21} useful muon decays per year.

(i) For $\sin^2 \theta_{13} \gtrsim 10^{-2}$, mass hierarchy can be determined to 5σ at almost all the baselines > 1000 km. For lower θ_{13} , baselines ~ 2500 km achieve the task for the largest fraction of δ_{CP} , however even they may fall short when δ_{CP} is near $3\pi/2$.

(ii) Any baseline $\lesssim 2000$ km can discover θ_{13} if $\sin^2 \theta_{13} \gtrsim 10^{-3}$. For smaller θ_{13} , the choice of optimal baseline depend on the actual value of δ_{CP} . However for majority of values of δ_{CP} the baseline range 1300 - 2500 km can be termed as optimal.

(iii) The discovery of δ_{CP} is naturally harder when its value is near 0 or π . However if $\sin^2 \theta_{13} \gtrsim 0.01$, the discovery is possible for a wide range of δ_{CP} values centered at $\pi/2$ and $3\pi/2$. The range typically decreases with the

increase of baseline.

(iv) At a given baseline, when the parent muon energy is increased, the performance typically increases up to some energy and then remains the same. We term this as the saturation behavior.

As can be gathered from above, there does not exist a unique “optimal” baseline or muon energy for all performance indicators. The baseline determination depends, in addition to scientific merit, also on geography, economics and sociology. Once that is determined, the optimization of muon energy involves mainly scientific and technical considerations. The saturation behavior mentioned above indicates that, given a baseline, it is preferable to have the parent muon energy as close to the saturation energy as possible, in order to avoid increasing the parent muon energy unnecessarily. While the saturation energy may be determined through a detailed simulation for a given baseline (as we have done for three benchmark baselines in this article), we have tried to come up with a simple criterion that can motivate this behavior and determine the saturation energy. We conjecture that the most efficient energy for a given baseline simply depends on the number of wrong-sign muons at the detector. This gives us a range of optimal muon energies for a given baseline, the range depending on the actual values of mixing parameters. This conjecture is vindicated a posteriori by the saturation energies obtained at the benchmark baselines. This simple-minded analysis gives an intuitive understanding of the relationship between baseline and the optimal energy.

With the recent indications of a large value of θ_{13} , the 5σ measurement of this quantity may already be within our grasp with the current experiments. Then the neutrino factory experiments have to aim higher, and the question to ask is what kind of setup will give us the 5σ determination of mass hierarchy and CP violation with the least amount of exposure needed. Our analysis indicates that for the mass hierarchy, the baseline of ~ 2500

km would perform the best at any parent muon energy. On the other hand, the measurement of CP violation is the most efficient around a baseline of ~ 1500 km at all parent muon energies, for most of the δ_{CP} values.

A few comments are in order. We have presented all our results assuming the actual hierarchy to be NH. However note that the probabilities obtained with NH and μ^+ beam are identical with the probabilities with IH and μ^- beam, and we have taken beams of both polarities with equal exposure. In LENF, both the polarities of muons can be accelerated at the same time, giving alternate bunches of μ^+ and μ^- . The cross sections of ν_μ and $\bar{\nu}_\mu$ on the nucleons are also virtually identical at the relevant energies, and hence our results are valid even for IH.

Another comment is about the measurement of δ_{CP} itself. In this article, we have only considered the discovery potential for CP violation, i.e. we are interested in finding a nonzero δ_{CP} . This task is, naturally, hard for small δ_{CP} and impossible when $\delta_{CP}=0$ or π . However with large θ_{13} , a measurement of δ_{CP} may be possible even if its value is close to the CP-conserving limit. If indeed the value of θ_{13} is confirmed to be large, one can aim to answer more detailed questions like the value of δ_{CP} , the octant of θ_{23} , or the deviation of θ_{23} from maximality. These quantities will not be analyzed in this work.

Finally, the most important quantity is the one we have no control over: the value of θ_{13} . As far as neutrino factories are concerned, this quantity will determine if they are going to be discovery machines or precision machines.

Acknowledgments

We thank S. Prakash, S. Raut and S. UmaSankar for useful discussions. S.R. would like to thank Yuval Grossman for support and hospitality.

-
- [1] G. L. Fogli, E. Lisi, A. Marrone, A. Palazzo and A. M. Rotunno, arXiv:1106.6028 [hep-ph].
 - [2] T. Schwetz, M. Tortola and J. W. F. Valle, arXiv:1108.1376 [hep-ph].
 - [3] K. Abe *et al.* [T2K Collaboration], arXiv:1106.2822 [hep-ex].
 - [4] L. Whitehead [MINOS Collaboration], “Recent results from MINOS” Fermilab Seminar, 24 June, 2011 http://www.numi.fnal.gov/pr_pLOTS/.
 - [5] A. de Gouvea, J. Jenkins and B. Kayser, Phys. Rev. D **71**, 113009 (2005) [arXiv:hep-ph/0503079]; A. de Gouvea and W. Winter, Phys. Rev. D **73**, 033003 (2006) [arXiv:hep-ph/0509359]; R. Gandhi, P. Ghoshal, S. Goswami and S. U. Sankar, Mod. Phys. Lett. A **25**, 2255 (2010) [arXiv:0905.2382 [hep-ph]].
 - [6] F. Ardellier *et al.*, arXiv:hep-ex/0405032; X. Guo *et al.* [Daya-Bay Collaboration], arXiv:hep-ex/0701029; J. K. Ahn *et al.* [RENO Collaboration], arXiv:1003.1391 [hep-ex].
 - [7] D. Indumathi and M. V. N. Murthy, Phys. Rev. D **71**, 013001 (2005) [arXiv:hep-ph/0407336]; R. Gandhi, P. Ghoshal, S. Goswami, P. Mehta and S. Uma Sankar, Phys. Rev. D **73**, 053001 (2006) [arXiv:hep-ph/0411252]; S. T. Petcov and T. Schwetz, Nucl. Phys. B **740**, 1 (2006) [arXiv:hep-ph/0511277]; R. Gandhi, P. Ghoshal, S. Goswami, P. Mehta, S. U. Sankar and S. Shalgar, Phys. Rev. D **76**, 073012 (2007) [arXiv:0707.1723 [hep-ph]]; A. Samanta, Phys. Lett. B **673**, 37 (2009) [arXiv:hep-ph/0610196].
 - [8] P. Huber, M. Lindner, T. Schwetz and W. Winter, JHEP **0911**, 044 (2009) [arXiv:0907.1896 [hep-ph]].
 - [9] A. Bandyopadhyay *et al.* [ISS Physics Working Group Collaboration], Rept. Prog. Phys. **72**, 106201 (2009), [arXiv:0710.4947 [hep-ph]].
 - [10] V. Barger, D. Marfatia and K. Whisnant, Phys. Rev. D **65**, 073023 (2002) [arXiv:hep-ph/0112119]; P. Hu-

- ber and W. Winter, Phys. Rev. D **68**, 037301 (2003) [arXiv:hep-ph/0301257].
- [11] <https://www.ids-nf.org>.
- [12] S. Geer, O. Mena and S. Pascoli, Phys. Rev. D **75**, 093001 (2007) [arXiv:hep-ph/0701258]; E. Fernandez Martinez, T. Li, S. Pascoli and O. Mena, Phys. Rev. D **81**, 073010 (2010) [arXiv:0911.3776 [hep-ph]].
- [13] S. K. Agarwalla, P. Huber, J. Tang and W. Winter, JHEP **1101**, 120 (2011) [arXiv:1012.1872 [hep-ph]].
- [14] P. Huber and T. Schwetz, Phys. Lett. B **669**, 294 (2008) [arXiv:0805.2019 [hep-ph]].
- [15] S. K. Raut, R. S. Singh, S. U. Sankar, Phys. Lett. **B696**, 227-231 (2011) [arXiv:0908.3741 [hep-ph]].
- [16] A. Dighe, S. Goswami and S. Ray, Phys. Rev. Lett. **105**, 261802 (2010) [arXiv:1009.1093 [hep-ph]].
- [17] M. V. Diwan *et al.*, Phys. Rev. D **68**, 012002 (2003) [arXiv:hep-ph/0303081].
- [18] J. Peltoniemi, arXiv:0911.4876 [hep-ex].
- [19] A. Joglekar, S. Prakash, S. K. Raut, S. U. Sankar, Mod. Phys. Lett. **A26**, 2051-2063 (2011) [arXiv:1011.1146 [hep-ph]]; S. K. Agarwalla, T. Li, A. Rubbia, [arXiv:1109.6526 [hep-ph]].
- [20] A. Y. Smirnov, arXiv:hep-ph/0610198.
- [21] E. K. Akhmedov, R. Johansson, M. Lindner, T. Ohlsson and T. Schwetz, JHEP **0404**, 078 (2004) [arXiv:hep-ph/0402175].
- [22] J. Burguet-Castell, M. B. Gavela, J. J. Gomez-Cadenas, P. Hernandez and O. Mena, Nucl. Phys. B **608**, 301 (2001); H. Minakata and H. Nunokawa, JHEP **0110**, 001 (2001).
- [23] G. L. Fogli and E. Lisi, Phys. Rev. D **54**, 3667 (1996);
- [24] A. M. Dziewonski and D. L. Anderson, Phys. Earth Planet. Inter. **25**, 297 (1981).
- [25] P. Huber and T. Schwetz, Phys. Lett. B **669**, 294 (2008) [arXiv:0805.2019 [hep-ph]].
- [26] E. Fernandez Martinez, T. Li, S. Pascoli and O. Mena, Phys. Rev. D **81**, 073010 (2010) [arXiv:0911.3776 [hep-ph]].
- [27] P. Huber, M. Lindner and W. Winter, Comput. Phys. Commun. **167**, 195 (2005) [arXiv:hep-ph/0407333]; P. Huber, J. Kopp, M. Lindner, M. Rolinec and W. Winter, Comput. Phys. Commun. **177**, 432 (2007) [arXiv:hep-ph/0701187].